

Wandering Seminar
Mini-conference on Ergodic Theory & Dynamical Systems
Jagiellonian University in Krakow, 24–26 January 2014

Abstracts of talks

January 20, 2014

Mini-course, Friday–Sunday, January 24–26

Dimension of self-similar sets with overlaps

Michael Hochman (Hebrew University of Jerusalem)

The lecture series will present recent work on the dimension of self similar sets and measures with overlaps, and some applications. Much of the analysis relies on ideas from additive combinatoric, and we will discuss these as well. The lectures are aimed at graduate students and researchers and will assume graduate-level background in real analysis and measure theory. Some familiarity with Minkowski dimension, Hausdorff dimension and Shannon entropy is recommended.

Friday, January 24

Multi-fractal spectra and their computation

Mark Pollicott (University of Warwick)

By Multifractal Analysis we mean the dimension of the set of points for which the cesaro averages of a given function along orbits of a transformation converge to different values. We will consider the special case of an expanding interval map and recall the approach based on thermodynamical formalism. Finally we will relate this to periodic points and determinants, which gives a useful approach to numerically computing these dimensions.

Fourier transforms of measures supported on graphs.

Jonathan Fraser (University of Warwick)

I will discuss the Fourier transforms of measures supported on graphs of real-valued functions defined on the unit interval. In particular, I will be interested in how fast these Fourier transforms converge to zero (if they do at all), which is related to the concept of Fourier dimension. Our results imply that the graph of fractional Brownian motion is almost surely not a Salem set, answering in part a question of Kahane from 1993, and that the graph of a Baire typical function in $C[0,1]$ has Fourier dimension zero.

This is joint work with Tuomas Orponen and Tuomas Sahlsten.

Hyperbolic dynamics in graph-directed IFS

Jacek Tabor (Jagiellonian University in Kraków)

In this talk our aim is to present an extension of the classical theory of fractal attractors build by contracting IFS in metric spaces to the hyperbolic case. As a trivial application of our method we obtain for example the estimations for the fractal dimension of the Smale's Horseshoe defined by a small Lipschitz perturbation of the classical Smale's map.

To construct our approach we need the notion of a *cone space*, which is a complete metric space (X, d) with a pair of functions $c_s, c_u : X \times X \rightarrow \mathbb{R}$, such that there exists $K > 0$ satisfying

$$\frac{1}{K}d(x, x') \leq \max(c_s(x, x'), c_u(x, x')) \leq Kd(x, x') \text{ for } x, x' \in X.$$

Cone space allows us to use the ideas standing behind the standard cone condition: for a partial map f between cone spaces X and Y we introduce $|f|_s$ which measures the stable contraction rate and $\langle f \rangle_u$ which measures the unstable expansion rate. We say that f is *cone-hyperbolic* if

$$|f|_s < 1 < \langle f \rangle_u.$$

Using cone hyperbolicity and graph-directed IFS we build an abstract metric model which describes the dynamics of the hyperbolic-like systems.

This allows to obtain estimations from below and above of the fractal dimension of the hyperbolic invariant set, which are exact analogs of the well-known classical results obtained for contracting IFS.

This is a joint work with Tomasz Kułaga. The results presented in our talk can be find in T. Kułaga, J. Tabor, *Hyperbolic dynamics in graph-directed IFS*, Journal of Differential Equations **251** (12), pp. 3363–3380.

Saturday, January 25

On the dimension of the graph of the classical Weierstrass functions

Part 1 — Transversality method

Balázs Bárány (Technical University of Budapest)

Part 2 — Ledrappier-Young Theory

Julia Romanowska (IM PAN Warszawa)

We examine dimension of the graph of the famous Weierstrass non-differentiable function. We prove that there exist sets A and B of parameters such that the Hausdorff dimension of the graph of the Weierstrass function is equal to its Lyapunov dimension for every parameters in A and Lebesgue almost every parameters in B . Furthermore, we prove that the Hausdorff dimension of the graph is equal to the Lyapunov dimension for typical \mathbb{Z} -periodic C^3 functions.

On examples of completely scrambled systems

Piotr Oprocha (AGH University of Science and Technology)

A dynamical system (X, f) is completely scrambled if every pair x, y of distinct points in X is proximal (i.e. $\liminf d(f^n(x), f^n(y)) = 0$) but not asymptotic (i.e. $\limsup d(f^n(x), f^n(y)) > 0$).

In 1999, Huang and Ye proved that continua of arbitrary (topological) dimension admit completely scrambled homeomorphisms. In this talk we aim to present a brief characterization of completely scrambled homeomorphisms, ideas behind the construction of Huang and Ye and some other methods of construction of completely scrambled systems which are, in fact, motivated by much earlier papers (on different topics).

If time permits, we will present a method to answer a question left open in the Huang and Ye's paper (joint work with Foryś, Huang and Li). Finally, we will present a few questions for further research.

Sunday, January 26

Lyapunov-optimizing measures for some matrix-valued cocycles

Michał Rams (IM PAN Warszawa)

Choose some finite set $A = \{A_1, \dots, A_k\}$ of 2×2 real-valued matrices. Consider the products $B(\omega_1, \dots, \omega_n) = A_{\omega_n} \cdot \dots \cdot A_{\omega_1}$ of the matrices belonging to this set, in any order, possibly using some matrices more than once. Try to make the norm of the product grow as fast as possible (that is, consider the function

$$f(\omega) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \|B(\omega_1, \dots, \omega_n)\|$$

defined on $\Sigma = \{1, \dots, k\}^{\mathbb{N}}$ and maximize it). Or try to have the norm grow as slowly as possible. How many possible choices of ω you have?

The intuition is that, under some not-too-restrictive assumptions (like none of the matrices appearing twice), there shouldn't be many growth-maximizing (or growth-minimizing) sequences. While this intuition is a little misleading, especially in the case of growth-minimizing sequences, we are able to show an open family of sets A for which all the shift-invariant measures on Σ maximizing/minimizing the norm growth have zero entropy. This is a joint work with Jairo Bochi.